

Biologia Geral e Experimental

Universidade Federal de Sergipe

São Cristóvão, SE 5 (2): 5-9

30.vii.2005

CARDIOVASCULAR EFFECTS ON RATS INDUCED BY THE TOTAL ALKALOID FRACTION OF *SIDA CORDIFOLIA*

Márcio Roberto Viana Santos¹

Murilo Marchioro¹

Aletéia Lacerda Silveira²

José Maria Barbosa Filho²

Isac Almeida Medeiros²

ABSTRACT

Cardiovascular effects induced by the total alkaloid fraction of *Sida cordifolia* L. (TAF) were investigated in unanesthetized normotensive rats. In these animals, TAF (0.5, 1, 5 and 10 mg/kg, i.v.) induced hypotension, which was attenuated by atropine, L-NAME or hexamethonium at the 10 mg/kg dose, and bradycardia, which was eliminated by atropine at all doses, and attenuated by hexamethonium at 5 and 10 mg/kg, but was unchanged by L-NAME. These results demonstrate that TAF induces hypotension and bradycardia, which appear to be mediated by direct and indirect activation of muscarinic receptors. Nitric oxide also appears to be involved in the hypotensive response.

Keywords: *Sida cordifolia*, total alkaloid fraction, cardiovascular effects, hypotension, rat.

RESUMO

Os efeitos cardiovasculares induzidos pela fração de alcalóides totais da *Sida cordifolia* L. (TAF) foram avaliados em ratos normotensos não-anestesiados. Nestes animais, TAF (0,5; 1; 5 e 10 mg/kg, i.v.) induziu hipotensão, que foi atenuada pela atropina, L-NAME ou hexametônio apenas na dose de 10 mg/kg, e bradicardia que foi abolida pela atropina, em todas as doses, e atenuada pelo hexametônio nas doses de 5 e 10 mg/kg, mas não foi alterada pelo L-NAME. Estes resultados demonstram que TAF induz hipotensão e bradicardia, que parece envolver uma ativação direta e indireta de receptores muscarínicos. Além disso, o óxido nítrico parece também participar do efeito hipotensor induzido pelo TAF.

Palavras-chave: *Sida cordifolia*, fração de alcalóides totais, efeitos cardiovasculares, hipotensão, rato.

INTRODUCTION

The use of medicinal plants for the treatment of human diseases has increased considerably throughout the World. Evaluation of the effects of these plants on organs and systems contributes to the development of the scientific basis for therapeutic applications, and enriches alternatives for the treatment of a growing number of diseases (Elizabetsky, 1986).

Sida cordifolia L. (Malvaceae), a native species

of the Brazilian Northeast, is known popularly as “Malva branca”. It is used in folk medicine as an antirheumatic and antipyretic (Muzaffer *et al.*, 1991), an antiinflammatory and analgesic (Franzotti *et al.*, 2000), and an antiasthmatic and nasal anticongestant (Ghosh & Dutt, 1930; Mukerji, 1953). Previous studies have demonstrated that vasicine, an alkaloid isolated from this plant (Ghosal *et al.*, 1975), produced hypotension and bradycardia in anaesthetized dogs (Gupta *et al.*, 1977) and unanaesthetized rats (Silveira

¹ Departamento de Fisiologia, Universidade Federal de Sergipe, Av. Marechal Rondon, s/n, Jardim Rosa Elze, São Cristóvão, Se, 49100-000. marcio@infonet.com.br

² Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, caixa postal 5009, João Pessoa, PB, 58.051-970.

et al., 2003). In the pharmacological study reported here, we evaluated the effects of the total alkaloid fraction of *Sida cordifolia* L. on the arterial pressure and heart rate of unanesthetized, normotensive rats (*Rattus norvegicus*). We demonstrate that the hydroalcoholic extract of this plant also induces marked hypotension, associated with intense bradycardia, in rats.

MATERIALS AND METHODS

Preparation of the total alkaloid fraction: *Sida cordifolia* leaves (voucher specimen n°. 30171, deposited in the herbarium of the Department of Biology, Universidade Federal de Sergipe, Brazil) were dried, pulverized and immersed in 95% EtOH at room temperature for 72 h. After vacuum concentration, the extract obtained was submitted to conventional acid-base treatment to produce the total alkaloid fraction (TAF). To avoid decomposition, the TAF was dissolved in a cremophor-saline solution (0.1% v/v) at desired concentrations just before each experiment. Cremophor had no effect when tested under controlled conditions (data not shown).

Animals: Male Wistar rats (250-350 g) were used for all experiments. The animals were housed under controlled conditions of temperature ($21 \pm 1^\circ\text{C}$) and lighting (lights on: 06:00-18:00 h), with free access to food and tap water.

Drugs: The drugs used were: heparin sodium salt (Ariston), sodium thiopental (Cristália), atropine sulfate, N^w-nitro-*L*-arginine methyl ester (L-NAME), hexamethonium, sodium nitroprusside and cremophor (all from Sigma). All drugs were dissolved freely in saline solution.

Blood pressure measurements: Under sodium thiopental anesthesia (45 mg/kg, i.v.), the lower

abdominal aorta and inferior *vena cava* were cannulated via the left femoral artery and vein using polyethylene catheters. The catheters were then filled with heparinized saline solution and slid under the skin to emerge between the scapulae. Arterial pressure was measured after 24 h by connecting the arterial catheter to a pre-calibrated pressure transducer (Edwards Lifescience, Irvine, CA, USA) and pressure outputs were recorded by an amplifier-recorder (BioData, Model BD-01, PB, Brazil) connected to a personal computer equipped with an analog-to-digital converter board (BioData, PB, Brazil). Data were collected at a frequency of 200 Hz. For each cardiac cycle, the computer calculated mean arterial pressure (MAP) and pulse interval, referred to here as the heart rate (HR). The venous catheter was inserted for drug administration. Sodium nitroprusside (10 µg/kg) was injected to check the efficacy of catheter insertion.

Experimental protocol: After stabilization of cardiovascular parameters, MAP and HR were recorded before (baseline values) and after randomized i.v. administration of different doses of TAF (0.5, 1, 5 and 10 mg/kg). These administrations were separated by a time interval sufficient to allow full recovery of cardiovascular parameters. A control dose-response curve was then obtained. Dose-response curve were also obtained separately in animals pre-treated with atropine (2 mg/kg; i.v.; 15 min.), a non-selective antagonist of muscarinic receptors (Mitchelson, 1984), L-NAME (20 mg/kg, i.v. 30 min.), a competitive inhibitor of NO-synthase (Moncada & Higgs, 1993) or hexamethonium, a ganglionic blockade (Takahashi & Owyang, 1997).

Statistics: Values are expressed as means \pm SEM. The significance of differences among means was tested using one-way ANOVA with Dunnett's post test. All procedures were carried out on Graph Pad PrismTM version 3.02 software.

RESULTS

As expected from control animals, atropine and hexamethonium increased HR significantly, from 350 ± 7 bpm to 485 ± 11 and 416 ± 20 bpm ($p < 0.05$; $n = 6$), respectively. The administration of L-NAME increased MAP and HR from 110 ± 3 mmHg to 136 ± 4 mmHg and from 350 ± 7 bpm to 373 ± 9 bpm ($p < 0.05$; $n = 6$), respectively.

Figure 1 shows original traces of the effect induced by *S. cordifolia* TAF (10 mg/kg, i.v.) in one unanesthetized normotensive rat. In these animals, TAF (0.5, 1, 5 and 10 mg/kg; i.v., randomly) induced transitory hypotension (-1 ± 1 ; -5 ± 1 ; -21 ± 4 and -39 ± 3 %) associated with intense bradycardia (-0.7 ± 0.1 ; -5 ± 0.7 ; -44 ± 7 and -70 ± 2 %) (Figure 2).

In animals pre-treated with atropine (2 mg/kg), L-NAME (20 mg/kg) or hexamethonium (20 mg/kg), the hypotensive response induced by TAF was only attenuated at the highest dose (10 mg/kg). However, the bradycardic response was eliminated by atropine at all doses, and attenuated by hexamethonium at 5 and 10 mg/kg, although it was not altered by L-NAME (Figure 2).

DISCUSSION

We chose to evaluate the effects of the TAF of *S. cordifolia* on the cardiovascular parameters of unanesthetized rats in order to avoid the possible influence of anesthesia and surgical stress (Smith & Hutchins, 1980; Fluckiger *et al.*, 1985). Baseline MAP and HR values recorded here were similar to those reported in previous studies (Lahlou *et al.*, 2002; Silveira *et al.*, 2003; Cunha *et al.*, 2004). Acute administration of TAF induced hypotension associated with bradycardia.

Peterson *et al.* (1984) established that the primary autonomic regulation of sinoatrial node function is carried out by vagal action via stimulation

of cardiac muscarinic receptors. Stimulation of these receptors induces intense bradycardia followed by hypotension due to a decrease in cardiac output. It is also well known that activation of endothelial muscarinic receptors induces intense vasodilatation due to release of endothelium-derived relaxing factors (Moncada *et al.*, 1991), mainly NO (Furchgott & Zawadzki, 1980; Moncada *et al.*, 1991). This activation can cause decrease in peripheral vascular resistance and, consequently, hypotension.

In order to evaluate the role of these receptors in TAF-induced responses, we performed experiments in the presence of atropine, a non-selective antagonist of muscarinic receptors (Mitchelson, 1984). Under these conditions, bradycardia was abolished at all doses, although hypotension was attenuated significantly at 10 mg/kg. This indicates that TAF induces bradycardia either through direct activation of receptors, or indirectly via vagal stimulation and release of acetylcholine into the sinoatrial node. However, hypotension does not appear to be due exclusively to the decrease in cardiac output related to bradycardia, given that it was only attenuated, but not abolished by atropine. This indicates that a decrease in peripheral vascular resistance may also contribute to TAF-induced hypotension.

To verify a possible indirect effect of TAF via the vagal nerve, we performed experiments with hexamethonium, a ganglionic blocker (Takahashi & Owyang, 1997). This drug attenuated bradycardia and hypotension significantly, but did not eliminate them altogether. This indicates that the bradycardic response is mediated by two distinct pathways, indirect and direct activation of cardiac muscarinic receptors.

Finally, we investigated the role of nitric oxide. In this condition, L-NAME, a competitive inhibitor of NO-synthase (Moncada & Higgs, 1993), changed the hypotensive response only at the 10 mg/kg dose, suggesting that NO is at least partly involved in the hypotensive effect.

These results agree with those of Silveira *et al.*




Figure 1. Original traces showing the effect of TAF (10 mg/kg; i.v.) on MAP and HR in one non-anaesthetized normotensive rat. The arrow indicates point of TAF administration.




Figure 2. Hypotensive and bradycardic response induced by TAF (0.5, 1, 5 and 10 mg/Kg; i.v.) in non-anaesthetized normotensive rats before (Control) and after pre-treatment with atropine (2 mg/kg), hexamethonium (20 mg/kg) or L-NAME (20 mg/kg). Values are expressed as mean \pm SEM of six experiments. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ vs Control.

(2003), who demonstrated that vasicine, an alkaloid isolated from *S. cordifolia* (Ghosal *et al.*, 1975), also produced marked hypotension associated with bradycardia in unanesthetized rats. The effects recorded in the present study may thus be due to the presence of vasicine in the TAF, although further data would be required to confirm this.

Overall, our results demonstrate that the TAF of *S. cordifolia* induces hypotension and bradycardia, apparently mediated by direct and indirect activation of muscarinic receptors. Nitric oxide also appears to be involved in the hypotensive response.

Acknowledgements: This study was supported by grants from FAP-SE and CAPES, Brazil.

REFERENCES

- Cunha, R.M., S.R.Q. Farias, J.C. Duarte, M.R.V. Santos, E.A.N. Ribeiro & I.A. Medeiros, 2004. Cardiovascular effects induced in rats by the essential oil of *Ocotea duckei* Vattimo (LAURACEAE), **Biol. Geral e Exp.** 5(1):12-18.
- Elizabetzky, E. 1986. New directions in ethnopharmacology. **J. Etnobiol.** 6:121.
- Fluckiger, J.P., M. Sonnay, N. Boillat, & J. Atkinson, 1985. Attenuation of the baroreceptor reflex by general anesthetic agent in the normotensive rat. **Eur. J. Pharmacol.** 109:105-109.
- Franzotti, E.M., C.V.F. Santos, H.M.S.L. Rodrigues, R.H.V. Mourão, M.R. Andrade & A.R. Antonioli, 2000. Anti-inflammatory, analgesic activity and acute toxicity of *Sida cordifolia* L. (Malva-Branca). **J. Ethnopharmacol.** 72(1/2):273-278.
- Furchgott, R.F. & J.V. Zawadzki, 1980. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. **Nature** 288: 373-376.
- Ghosal, S., R.B.P.S. Chauhan & R. Mehta, 1975. Chemical constituents of Malvaceae. Part I. Alkaloids of *Sida cordifolia*. **Phytochemistry** 14:830-832.
- Ghosh, S. & A. Dutt, 1930. The chemical examination of *Sida cordifolia*. **J. Indian Chem. Soc.** 7:825.
- Gupta, O.P., M.L. Sharma, B.J. Ray Ghatak & C.K. Atal, 1977. Pharmacological investigations of vasicine and vasicinone – the alkaloids of *Adhatoda vasica*. **Indian J. Med. Res.** 66: 680-691.
- Lahlou S, Figueiredo AF, Magalhaes PJ, Leal-Cardoso JH (2002) Cardiovascular effects of 1,8-cineole, a terpenoid oxide present in many plant essential oils, in normotensive rats. **Can. J. Physiol. Pharmacol.** 80(12):1125-1131.
- Mitchelson, F. 1984. Heterogeneity in muscarinic receptors: evidence from pharmacological studies with antagonists. **Trends Pharmacol. Sci. (suppl.)** 5:12-16.
- Moncada, S. & E.A. Higgs, 1993. The L-arginine-nitric oxide pathway. **New England Journal of Medicine** 29:2002-2012.
- Moncada, S., R.M.J. Palmer & E.A. Higgs, 1991. Nitric Oxide: Physiology, pathophysiology and pharmacology. **Pharmacol. Rev.** 43:109-142.
- Mukerji, B. 1953. The Indian Pharmaceutical Codex. Volume I - Indigenous Drugs. **Council of Scientific and Industrial Research**, New Delhi, India.
- Muzaffer, A.; S. Joy & S. Usmanu Ah, 1991. Screening of *Sida cordifolia* Linn, *Sida rhomboidea* Linn and *Triumfetta rotundifolia* Lam. for anti-inflammatory and anti-pyretic activities. **Indian Drugs** 28(9): 397.
- Peterson, G.L., G.S. Herron, M. Yamaki, D.S. Fullerton & M.I. Schimerlik, 1984. Purification of the muscarinic acetylcholine receptor from porcine atria. **Proc. Natl. Acad. Sci. U S A** 81:4993-4997.
- Smith, T.L. & P.M. Hutchins, 1980. Anesthetic effects on hemodynamics of spontaneously hypertensive and Wistar-Kyoto rats. **Am. J. Physiol.** 238: H539-H544.
- Takahashi, T. & C. Owyang, 1997. Characterization of vagal pathways mediating gastric accommodation reflex in rats. **J. Physiol.** 504(2):479-488.
- Silveira, A.L, M.A.S. Gomes, R.N. Silva Filho, M.R.V. Santos, I.A. Medeiros & J.M. Barbosa-Filho, 2003. Evaluation of the cardiovascular effects of vasicine, an alkaloid isolated from the leaves of *Sida cordifolia* L. (MALVACEAE). **Rev. Bras. Farmacognosia** 13(supl 2):37-39.

Biologia Geral e Experimental

Universidade Federal de Sergipe

São Cristóvão, SE 5 (2): 10-15

30.vii.2005

ANTINOCICEPTIVE ACTIVITY OF THE AQUEOUS EXTRACT OF *BAUHINIA CHEILANTHA* (BONG.) STEUD. (LEGUMINOSAE: CAESALPINIOIDEAE)

Adriana M. O. Silva¹
Flavia Teixeira-Silva¹
Rogéria de Souza Nunes¹
Rosilene Moretti Marçal¹
Sócrates Cabral de Holanda Cavalcanti¹
Ângelo Roberto Antonioli¹

ABSTRACT

Bauhinia cheilantha (Leguminosae: Caesalpinioideae) is a common plant of the Brazilian Caatinga widely used in folk medicine as an analgesic. In order to verify this effect pharmacologically, the antinociceptive activity of the plant was evaluated through the administration of its aqueous extract in mice. The extract was administered orally (400 mg/kg) 60 minutes before a writhing test, and was found to reduce nociception by 54.4%. The effects of formalin (1%) were also reduced by the extract at all doses. Naloxone (5 mg/kg, i.p.) and caffeine (10 mg/kg, i.p) reverted the effect of the extract. In a hot plate test, the extract (100mg, 200mg and 400 mg/kg) increased latency time by 39.8%, 30.7% and 32.8%, respectively. There was no acute toxicity in doses up to 3g/kg. The aqueous extract of the *B. cheilantha* bark revealed antinociceptive activity in all the models tested, effects that are possibly associated with the opioid and adenosine systems.

Keywords: *Bauhinia cheilantha*, antinociceptive, aqueous extract.

RESUMO

Bauhinia cheilantha (Leguminosae-Caesalpinioideae) é uma planta comum da caatinga amplamente utilizada na medicina popular como analgésica. Para verificar farmacologicamente este efeito, a atividade antinociceptiva desta planta foi verificada através do extrato aquoso da entre-casca, administrado em camundongos. O extrato (400 mg/kg), administrado por via oral 60 minutos antes do teste reduziu a nocicepção em 54,4%. Os efeitos da formalina (1%) também foram reduzidos pelo extrato em todas as doses. Naloxona (5 mg/kg, i.p.) e cafeína (10 mg/kg, i.p) também reverteram este efeito. No teste da placa quente o extrato de *B. cheilantha* nas doses 100mg, 200mg e 400 mg/kg, aumentou o tempo de latência em 39.8%, 30.7% e 32.8%, respectivamente. Não houve toxicidade aguda até a dose de 3g/Kg. O extrato aquoso da entre-casca de *B. cheilantha* mostrou atividade antinociceptiva nos modelos testados, efeito possivelmente associado com os sistemas opióide e adenosina.

Palavras-chave: *Bauhinia cheilantha*, antinociceptivo, extrato aquoso.

INTRODUCTION

Many plants of the Brazilian flora are used in local folk medicine, but very few have been pharmacologically validated. Among the unstudied majority is the legume *Bauhinia cheilantha* (Bong.) Steud., a small tree (average height 5 m), which is relatively common in the semi-arid Brazilian Caatinga.

Known locally as *mororó*, the extract of its bark is used as an analgesic, but there are no pharmacological data on its properties.

A variety of pharmacological properties have been reported for species of the genus *Bauhinia*. For example, the seeds of *Bauhinia monandra* affect haemagglutination positively, although this effect is eliminated by autoclaving (Abreu et al, 1990; Penate et

¹ Departamento de Fisiologia, Universidade Federal de Sergipe, Av. Marechal Rondon, s/n, Jardim Rosa Elze, São Cristóvão, Se, 49100-000. rosilenemoretti@ufl.br